



TABLE 111-3 Dosage Programs for Specific Conditions

| CONDITION   | FREQUENCY × DOSAGE  | TREATMENT PERIOD                                | COMBINATION* |
|---|---|---|--------------|
| Post-thrombotic syndrome <sup>209</sup>                               | 3 × 5   | As needed                                       | A            |
| Oncology <sup>210</sup>   | 2 × 11  | 6 wks or as needed                              | A            |
| Lymphedema <sup>141</sup>   | 2 × 5   | 2 yr  | A            |
| Fibrocystic breast disease <sup>177</sup>                             | 2 × 10  | 6 wks or as needed                              | A            |
| Experimental hematoma <sup>93</sup>                                   | 3 × 10  | 24 hr   | A            |
| Prophylactic treatment of karate injuries <sup>211</sup>              | 3 × 5   | During the season                               | A            |
| Traumatology and meniscus operation <sup>107</sup>                    | 3 × 10  | Average of 17.7 days                            | A            |
| Soft tissue injuries and distortions of the ankle joint <sup>94</sup> | 3 × 10 initially, then 3 × 1  | As needed                                       | A            |
| Athletic soft tissue injuries <sup>98</sup>                           | 2 × 5   | Spring football season                          | A            |
| Dental surgery <sup>115</sup>   | 4 × 5   | Day before surgery to seventh postoperative day | A            |
| Sports injuries <sup>101</sup>  | 3 × 10  | As needed                                       | A            |
| Chronic polyarthritis <sup>212</sup>                                  | 4 × 8 tablets   | 6 mo  | A            |
| Active arthroses <sup>213</sup>                                       | 4 × 7   | 6 wks   | A            |
| Extra-articular rheumatism <sup>144</sup>                             | 3 × 10 tablets  | As needed                                       | B            |
| Adnexitis <sup>127</sup>  | 3 × 5   | 10-14 days                                      | A            |
| Prostatitis <sup>117,125</sup>  | 3 × 5   | 10 days   | A            |
| Mastopathy <sup>214</sup>   | 2 × 10  |   | A            |
| Multiple sclerosis <sup>168-171,199</sup>                             |   |   |              |
| Initially   | 2 ampules injection   | Daily   | C            |
| As symptoms improve   | 2 ampules   | Every 2 to 3 days, then at longer intervals     | C            |
| On injection-free days  | 3 × 10  |   | A            |
| Later   | 3 × 5   |   | A            |
| Finally   | 3 × 3   |   | A            |
| Herpes Zoster <sup>185</sup>  |   |   |              |
| Initially   | 2 × 1 ampules   | First 2 days                                    | C            |
| First 2 days  | 3-4 tablets/day   | As indicated, up to 14 days                     | A            |
| HIV <sup>175,176</sup>  | 3 × 10 (ongoing studies insufficient to give effective dose levels) | As indicated                                    | A            |

\*For composition of combinations, see Table 111-2.